

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C11D 3/39, C07D 211/76		A1	(11) International Publication Number: WO 95/00626 (43) International Publication Date: 5 January 1995 (05.01.95)
(21) International Application Number: PCT/US94/06798 (22) International Filing Date: 16 June 1994 (16.06.94) (30) Priority Data: 08/082,207 24 June 1993 (24.06.93) US		(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors: WILLEY, Alan, David; 1071 Celestial Street, Cincinnati, OH 45202 (US). BURNS, Michael, Eugene; 9248 Sutherland Way, West Chester, OH 45069 (US). TSUNETSUGU, Shuichi; 1440 West Kemper Road #1805, Forest Park, OH 45240 (US). (74) Agents: REED, T., David, et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: BLEACHING COMPOUNDS COMPRISING ACYL VALEROLACTAM BLEACH ACTIVATORS

(57) Abstract

Laundry detergents and bleaching systems comprise acyl valerolactam bleach activators. The bleach activators are effective under mixed soil conditions, especially mixtures of hydrophobic and hydrophilic soils and stains.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MD	Republic of Moldova	TJ	Tajikistan
DE	Germany	MG	Madagascar	TT	Trinidad and Tobago
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain	MN	Mongolia	US	United States of America
FI	Finland			UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

1
BLEACHING COMPOUNDS COMPRISING ACYL
VALEROLACTAM BLEACH ACTIVATORS

5

FIELD OF THE INVENTION

10 The present invention relates to laundry detergents and bleaching systems which comprise acyl valerolactam bleach activators.

BACKGROUND OF THE INVENTION

15 It has long been known that peroxygen bleaches are effective for stain and/or soil removal from fabrics, but that such bleaches are temperature dependent. At a laundry liquor temperature of 60°C, peroxygen bleaches are only partially effective. As the laundry liquor temperature is lowered below 60°C, peroxygen bleaches become relatively ineffective. As a consequence, there has been a substantial amount of industrial research to develop bleaching systems which contain an activator that renders peroxygen bleaches effective at laundry liquor temperatures below 60°C.

20 Numerous substances have been disclosed in the art as effective bleach activators. One widely-used bleach activator is tetraacetyl ethylene diamine (TAED). TAED provides effective hydrophilic cleaning especially on beverage stains, but has limited performance on dingy stains and body soils. Another type of activator, such as nonanoyloxybenzenesulfonate (NOBS) and other activators which generally comprise long chain alkyl moieties, is hydrophobic in nature and provides excellent performance on dingy stains. However, many of the hydrophobic activators developed thus far can promote damage to natural rubber parts used in certain washing machines and to natural rubber articles exposed to the activators. Because of these negative effects on natural rubber machine parts and articles, the selection of such detergent-added
30 bleaching systems has been limited.

 It has now been determined that in conventional bleaching systems, particularly those comprising a hydrophobic bleach activator and a source of hydrogen peroxide, the bleach activator undergoes perhydrolysis to form a peroxyacid bleaching agent. A by-product of the perhydrolysis reaction between such bleach activators and hydrogen peroxide is a diacylperoxide (DAP) species. It has now further been discovered that the
35 DAP's derived from hydrophobic activators tend to be insoluble, poorly dispersible, oily materials which form a residue which can deposit on the natural rubber machine parts that are exposed to the laundry liquor. The oily DAP residue can form a film on the natural rubber parts and promote free radical and peroxide damage to the rubber, which

eventually leads to failure of the part. This is particularly true of rubber parts which have prolonged exposure to the laundry liquor, such as sump hoses.

By the present invention, it has now been discovered that the class of bleach activators derived from acyl valerolactams forms peroxyacids upon perhydrolysis without the production of oily, harmful DAP's. Without intending to be limited by theory, it is believed that the bleach activators employed herein provide good cleaning performance with safety to natural rubber, since they do not expose the natural rubber machine parts or articles to DAP oxidation. Whatever the reason, natural rubber parts and articles remain substantially undamaged by the bleaching systems of the present invention.


By the present invention, it has also now been discovered the bleach activators of this invention provide dingy soil clean-up and enhanced nucleophilic and body soil removal. Furthermore, the bleaching systems and activators herein are effective at low concentration levels and at temperatures below 60°C when used in the manner provided by this invention. In addition, the activators herein have better perhydrolysis speed and yield when compared to other lactam bleach activators, such as acyl caprolactam activators.

Accordingly, the present invention presents an effective, color-safe bleaching system which does not promote damage to natural rubber parts in washing machines or damage to natural rubber articles.

SUMMARY OF THE INVENTION


The present invention relates to acyl valerolactam bleach activators and their use in bleaching systems and laundry detergents. The valerolactams are selected from the group consisting of:

a)



wherein R¹ is a substituted or unsubstituted, including saturated or unsaturated, alkyl or alkoxy group containing from about 1 to about 18 carbon atoms wherein the longest linear alkyl or alkoxy chain extending from and including the carbonyl carbon contains from about 2 to about 12 carbon atoms;

b)



i.e., substituted and unsubstituted benzoyl valerolactams wherein substituents R², R³, R⁴, R⁵, and R⁶ are members selected from the group consisting of H, halogen, alkyl,

alkoxy, alkoxyaryl, alkaryl, and alkaryloxy moieties having from about 1 to about 12 carbon atoms, preferably from about 3 to about 12 carbon atoms, and substituents having the structure:



wherein R^7 is selected from the group consisting of H, alkyl, alkaryl, alkoxy, alkoxyaryl, alkaryloxy, and aminoalkyl; X is O, NH, or NR^9 , wherein R^9 is H or a C_1 - C_4 alkyl group; R^8 is an alkyl, cycloalkyl, or aryl group containing from 3 to 11 carbon atoms; and

c) mixtures of a) and b).


In a preferred embodiment of structure a), R^1 is selected from the group consisting of alkyl or alkoxy units having from about 7 to about 11 carbon atoms, including heptyl, octyl, nonyl, decyl, undecyl, decenyl, 2,4,4-trimethylpentyl, 1-ethylpentyl, and mixtures thereof.

In a preferred embodiment of structure b), R^2 , R^3 , R^4 , and R^5 are H and R^6 is selected from the group consisting of H, methyl, methoxy, ethyl, ethoxy, propyl, propoxy, isopropyl, isopropoxy, butyl, tert-butyl, butoxy, tert-butoxy, pentyl, pentoxy, hexyl, hexoxy, Cl, and NO_2 . In still another preferred embodiment, R^2 , R^3 , R^4 are H, and R^5 and R^6 are members selected from the group consisting of methyl, methoxy, and Cl.


The invention also relates to bleaching systems and laundry detergents comprising the bleach activators. Said bleaching system comprises:

A) at least about 0.1%, preferably from about 1% to about 75%, by weight of bleaching system, of a peroxygen bleaching compound capable of yielding hydrogen peroxide in an aqueous solution;

B) at least about 0.1%, preferably from about 0.1% to about 50%, by weight, of one or more acyl valerolactam bleach activators selected from the group consisting of:



wherein R^1 is as defined above;



wherein R^2 , R^3 , R^4 , R^5 , and R^6 are as defined above; and

c) mixtures of a) and b).

The peroxygen bleaching compound can be any peroxide source, and is preferably a member selected from the group consisting of sodium perborate monohydrate, sodium perborate tetrahydrate, sodium pyrophosphate peroxyhydrate, urea peroxyhydrate, sodium percarbonate, sodium peroxide and mixtures thereof.

- 5 Highly preferred peroxygen bleaching compounds are selected from the group consisting of sodium perborate monohydrate, sodium perborate tetrahydrate, sodium percarbonate and mixtures thereof. The most highly preferred peroxygen bleaching compound is sodium percarbonate.

- 10 The invention also encompasses laundry compositions in granular, paste, liquid, or bar form which comprise at least about 0.1% of the aforesaid bleaching system together with at least about 1% of conventional detergent ingredients which are present in the composition at the levels indicated hereinafter.

- The acyl valerolactams herein can also be used in combination with other bleach activators, such as N-acyl caprolactams, tetraacetyl ethylene diamine, 15 alkanoxybenzenesulfonates, including nonanoyloxybenzenesulfonate, benzoxazin-type bleach activators, and peroxyacid agents and activators having amide moieties. Preferably, if using an automatic washing machine equipped with natural rubber parts or if washing articles comprising natural rubber, the amount of alkanoxybenzenesulfonates used in combination with the activators of this invention should be 20 kept at a minimum.

- The bleaching method herein is preferably conducted with agitation of the fabrics with an aqueous liquor containing the aforesaid bleaching system at levels from about 50 ppm to about 27,500 ppm. The method can be carried out at any desired washing temperature, even at temperatures below about 60°C, and is readily conducted at 25 temperatures in the range of from about 5°C to about 45°C. The method can be conducted conveniently using a composition which is in bar form, but can also be conducted using granules, flakes, powders, pastes, liquids and the like.

- The aqueous laundry liquor typically comprises at least about 300 ppm of conventional detergent ingredients, as well as at least about 25 ppm of the bleaching 30 compound and at least about 25 ppm of the bleach activator. Preferably, the liquor comprises from about 900 ppm to about 20,000 ppm of conventional detergent ingredients, from about 100 ppm to about 25,000 ppm of the bleaching compound and from about 100 ppm to about 2,500 ppm of the bleach activator. The conventional detergent ingredients and bleaching system will typically be combined into a detergent 35 composition such as a granular laundry detergent or laundry detergent bar.

The conventional detergent ingredients employed in said method and in the compositions herein comprise from about 1% to about 99.8%, preferably from about 5% to about 80%, of a deterative surfactant. Optionally, the detergent ingredients

comprise from about 5% to about 80% of a detergent builder. Other optional detergent adjuncts can also be included in such compositions at conventional usage levels.

All percentages, ratios, and proportions herein are by weight, unless otherwise specified. All documents cited are incorporated herein by reference.

DETAILED DESCRIPTION OF THE INVENTION

5 The bleaching system employed in the present invention provides effective and efficient surface bleaching of fabrics which thereby removes stains and/or soils from the fabrics. The bleaching system is particularly efficient at cleaning concentrated soil loads, especially mixtures of hydrophobic and hydrophilic soils. Hydrophobic soils are
10 generally associated with nucleophilic, lipid and protein-based soils and stains, such as body soils, blood, etc., but are also effective on so-called "dingy soils". Dingy soils are those that build up on textiles after numerous cycles of usage and washing, and result in a gray or yellow tint on white fabrics. Hydrophilic soils include food and beverage stains. Further, the bleaching system is safe to natural rubber machine parts and
15 articles.

The bleaching mechanism and, in particular, the surface bleaching mechanism are not completely understood. However, it is generally believed that the bleach activator undergoes nucleophilic attack by a perhydroxide anion, which is generated from the hydrogen peroxide evolved by the peroxygen bleaching compound, to form a
20 peroxycarboxylic acid. This reaction is commonly referred to as perhydrolysis. It is also believed, that the bleach activators within this invention can render peroxygen bleaches more efficient even at laundry liquor temperatures wherein bleach activators are not necessary to activate the bleach, i.e., above about 60°C. Therefore, with bleach systems of the invention, less peroxygen bleach is required to achieve the same level of
25 surface bleaching performance as is obtained with the peroxygen bleach alone.

The components of the bleaching system herein comprise the bleach activator and the peroxide source, as described hereinafter.

Bleach Activators

Methods of making acyl valerolactams are illustrated by laboratory syntheses
30 included in Examples I and II.

Preferred valerolactams of structure a) include those wherein the R¹ moiety is selected from alkyl and alkoxy groups containing from about 7 to about 11 carbon atoms. Examples of preferred valerolactams include octanoyl valerolactam, nonanoyl valerolactam, decanoyl valerolactam, undecenoyl valerolactam, 3,5,5-trimethylhexanoyl
35 valerolactam, 2-ethylhexanoyl valerolactam, isononanoyl valerolactam and mixtures thereof. Highly preferred valerolactams of structure a) include nonanoyl valerolactam, isononanoyl valerolactam, and 2-ethylhexanoyl valerolactam.

Preferred valerolactams of structure b) include benzoyl valerolactam, substituted benzoyl valerolactam, including alkaryl and alkoxyaryl valerolactams wherein the alkaryl or alkoxyaryl moiety contains from about 3 to about 12 carbon atoms, terephthaloyl divalerolactam, and mixtures thereof. Examples of substituted benzoyl valerolactams include methylbenzoyl valerolactam, ethylbenzoyl valerolactam, ethoxybenzoyl valerolactam, propylbenzoyl valerolactam, propoxybenzoyl valerolactam, isopropylbenzoyl valerolactam, isopropoxybenzoyl valerolactam, butylbenzoyl valerolactam, butoxybenzoyl valerolactam, tertbutylbenzoyl valerolactam, tertbutoxybenzoyl valerolactam, pentylbenzoyl valerolactam, pentoxybenzoyl valerolactam, hexylbenzoyl valerolactam, hexoxybenzoyl valerolactam, 2,4,6-trichlorobenzoyl valerolactam, pentafluorobenzoyl valerolactam, dichlorobenzoyl valerolactam, dimethoxybenzoyl valerolactam, 4-nitrobenzoyl valerolactam, 3-chlorobenzoyl valerolactam, 4-chlorobenzoyl valerolactam, 2,4-dichlorobenzoyl valerolactam, terephthaloyl divalerolactam, and mixtures thereof. Highly preferred valerolactams of structure b) include benzoyl valerolactam, 3-chlorobenzoyl valerolactam, and 4-nitrobenzoyl valerolactam.

The bleaching system comprises at least about 0.1%, preferably from about 0.1% to about 50%, more preferably from about 1% to about 30%, most preferably from about 3% to about 25%, by weight, of one or more acyl valerolactam bleach activators. In highly preferred embodiments, the bleaching system comprises percarbonate and a bleach activator selected from the group consisting of benzoyl valerolactam, nonanoyl valerolactam, isononanoyl valerolactam, 2-ethylhexanoyl valerolactam, 3-chlorobenzoyl valerolactam, and 4-nitrobenzoyl valerolactam.

When the activators are used, optimum surface bleaching performance is obtained with washing solutions wherein the pH of such solution is between about 7 and 10.5, preferably between about 8.5 and 10.5, most preferably about 9.5 to about 10.5, in order to facilitate the perhydrolysis reaction. Such pH can be obtained with substances commonly known as buffering agents, which are optional components of the bleaching systems herein.

The Peroxygen Bleaching Compound

The peroxygen bleaching compounds useful herein are those capable of yielding hydrogen peroxide in an aqueous liquor. These compounds are well known in the art and include hydrogen peroxide and the alkali metal peroxides, organic peroxide bleaching compounds such as urea peroxide, and inorganic persalt bleaching compounds, such as the alkali metal perborates, percarbonates, perphosphates, and the like. Mixtures of two or more such bleaching compounds can also be used, if desired.

Preferred peroxygen bleaching compounds include sodium perborate, commercially available in the form of mono-, tri-, and tetra-hydrate, sodium

pyrophosphate peroxyhydrate, urea peroxyhydrate, sodium peroxide, and sodium percarbonate. Particularly preferred are sodium perborate tetrahydrate, sodium perborate monohydrate and sodium percarbonate. Sodium percarbonate is especially preferred because it is very stable during storage and yet still dissolves very quickly in the bleaching liquor. It is believed that such rapid dissolution results in the formation of higher levels of percarboxylic acid and, thus, enhanced surface bleaching performance.

Highly preferred percarbonate can be in uncoated or coated form. The average particle size of uncoated percarbonate ranges from about 400 to about 1200 microns, most preferably from about 400 to about 600 microns. If coated percarbonate is used, the preferred coating materials include mixtures of carbonate and sulphate, silicate, borosilicate, or fatty carboxylic acids.

The bleaching system comprises at least about 0.1%, preferably from about 1% to about 75%, more preferably from about 3% to about 40%, most preferably from about 3% to about 25%, by weight, of said peroxygen bleaching compound capable of yielding hydrogen peroxide in an aqueous solution.

The weight ratio of bleach activator to peroxygen bleaching compound in the bleaching system typically ranges from about 2:1 to 1:5. In preferred embodiments, the ratio ranges from about 1:1 to about 1:3.

The bleach activator/bleaching compound systems herein are useful per se as bleaches. However, such bleaching systems are especially useful in compositions which can comprise various detergent adjuncts such as surfactants, builders, enzymes, and the like as disclosed hereinafter. Such laundry detergent compositions comprise at least about 0.1%, preferably from about 1% to about 50%, of the bleaching system and at least about 1%, preferably from about 50% to about 99.9%, of conventional detergent ingredients.

Detergent Surfactant


The amount of detergent surfactant included in the fully formulated detergent compositions afforded by the present invention can vary from about 1% to about 99.8%, by weight of the detergent ingredients, depending upon the particular surfactants used and the effects desired. Preferably, the detergent surfactants comprise from about 5% to about 80%, by weight of the detergent ingredients.

The detergent surfactant can be nonionic, anionic, ampholytic, zwitterionic, or cationic. Mixtures of these surfactants can also be used. Preferred detergent compositions comprise anionic detergent surfactants or mixtures of anionic surfactants with other surfactants, especially nonionic surfactants.

Nonlimiting examples of surfactants useful herein include the conventional C₁₁-C₁₈ alkyl benzene sulfonates and primary, secondary, and random alkyl sulfates, the C₁₀-C₁₈ alkyl alkoxy sulfates, the C₁₀-C₁₈ alkyl polyglycosides and their corresponding

sulfated polyglycosides, C₁₂-C₁₈ alpha-sulfonated fatty acid esters, C₁₂-C₁₈ alkyl and alkyl phenol alkoxylates (especially ethoxylates and mixed ethoxy/propoxy), C₁₂-C₁₈ betaines and sulfobetaines ("sultaines"), C₁₀-C₁₈ amine oxides, and the like. Other conventional useful surfactants are listed in standard texts.

- 5 One particular class of adjunct nonionic surfactants especially useful herein comprises the polyhydroxy fatty acid amides of the formula:



- wherein: R¹ is H, C₁-C₈ hydrocarbyl, 2-hydroxyethyl, 2-hydroxypropyl, or a mixture thereof, preferably C₁-C₄ alkyl, more preferably C₁ or C₂ alkyl, most preferably C₁ alkyl (i.e., methyl); and R² is a C₅-C₃₂ hydrocarbyl moiety, preferably straight chain C₇-C₁₉ alkyl or alkenyl, more preferably straight chain C₉-C₁₇ alkyl or alkenyl, most preferably straight chain C₁₁-C₁₉ alkyl or alkenyl, or mixture thereof, and Z is a polyhydroxyhydrocarbyl moiety having a linear hydrocarbyl chain with at least 2 (in the case of glyceraldehyde) or at least 3 hydroxyls (in the case of other reducing sugars) directly connected to the chain, or an alkoxylated derivative (preferably ethoxylated or propoxylated) thereof. Z preferably will be derived from a reducing sugar in a reductive amination reaction; more preferably Z is a glycityl moiety. Suitable reducing sugars include glucose, fructose, maltose, lactose, galactose, mannose, and xylose, as well as
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995
 1000
 1005
 1010
 1015
 1020
 1025
 1030
 1035
 1040
 1045
 1050
 1055
 1060
 1065
 1070
 1075
 1080
 1085
 1090
 1095
 1100
 1105
 1110
 1115
 1120
 1125
 1130
 1135
 1140
 1145
 1150
 1155
 1160
 1165
 1170
 1175
 1180
 1185
 1190
 1195
 1200
 1205
 1210
 1215
 1220
 1225
 1230
 1235
 1240
 1245
 1250
 1255
 1260
 1265
 1270
 1275
 1280
 1285
 1290
 1295
 1300
 1305
 1310
 1315
 1320
 1325
 1330
 1335
 1340
 1345
 1350
 1355
 1360
 1365
 1370
 1375
 1380
 1385
 1390
 1395
 1400
 1405
 1410
 1415
 1420
 1425
 1430
 1435
 1440
 1445
 1450
 1455
 1460
 1465
 1470
 1475
 1480
 1485
 1490
 1495
 1500
 1505
 1510
 1515
 1520
 1525
 1530
 1535
 1540
 1545
 1550
 1555
 1560
 1565
 1570
 1575
 1580
 1585
 1590
 1595
 1600
 1605
 1610
 1615
 1620
 1625
 1630
 1635
 1640
 1645
 1650
 1655
 1660
 1665
 1670
 1675
 1680
 1685
 1690
 1695
 1700
 1705
 1710
 1715
 1720
 1725
 1730
 1735
 1740
 1745
 1750
 1755
 1760
 1765
 1770
 1775
 1780
 1785
 1790
 1795
 1800
 1805
 1810
 1815
 1820
 1825
 1830
 1835
 1840
 1845
 1850
 1855
 1860
 1865
 1870
 1875
 1880
 1885
 1890
 1895
 1900
 1905
 1910
 1915
 1920
 1925
 1930
 1935
 1940
 1945
 1950
 1955
 1960
 1965
 1970
 1975
 1980
 1985
 1990
 1995
 2000
 2005
 2010
 2015
 2020
 2025
 2030
 2035
 2040
 2045
 2050
 2055
 2060
 2065
 2070
 2075
 2080
 2085
 2090
 2095
 2100
 2105
 2110
 2115
 2120
 2125
 2130
 2135
 2140
 2145
 2150
 2155
 2160
 2165
 2170
 2175
 2180
 2185
 2190
 2195
 2200
 2205
 2210
 2215
 2220
 2225
 2230
 2235
 2240
 2245
 2250
 2255
 2260
 2265
 2270
 2275
 2280
 2285
 2290
 2295
 2300
 2305
 2310
 2315
 2320
 2325
 2330
 2335
 2340
 2345
 2350
 2355
 2360
 2365
 2370
 2375
 2380
 2385
 2390
 2395
 2400
 2405
 2410
 2415
 2420
 2425
 2430
 2435
 2440
 2445
 2450
 2455
 2460
 2465
 2470
 2475
 2480
 2485
 2490
 2495
 2500
 2505
 2510
 2515
 2520
 2525
 2530
 2535
 2540
 2545
 2550
 2555
 2560
 2565
 2570
 2575
 2580
 2585
 2590
 2595
 2600
 2605
 2610
 2615
 2620
 2625
 2630
 2635
 2640
 2645
 2650
 2655
 2660
 2665
 2670
 2675
 2680
 2685
 2690
 2695
 2700
 2705
 2710
 2715
 2720
 2725
 2730
 2735
 2740
 2745
 2750
 2755
 2760
 2765
 2770
 2775
 2780
 2785
 2790
 2795
 2800
 2805
 2810
 2815
 2820
 2825
 2830
 2835
 2840
 2845
 2850
 2855
 2860
 2865
 2870
 2875
 2880
 2885
 2890
 2895
 2900
 2905
 2910
 2915
 2920
 2925
 2930
 2935
 2940
 2945
 2950
 2955
 2960
 2965
 2970
 2975
 2980
 2985
 2990
 2995
 3000
 3005
 3010
 3015
 3020
 3025
 3030
 3035
 3040
 3045
 3050
 3055
 3060
 3065
 3070
 3075
 3080
 3085
 3090
 3095
 3100
 3105
 3110
 3115
 3120
 3125
 3130
 3135
 3140
 3145
 3150
 3155
 3160
 3165
 3170
 3175
 3180
 3185
 3190
 3195
 3200
 3205
 3210
 3215
 3220
 3225
 3230
 3235
 3240
 3245
 3250
 3255
 3260
 3265
 3270
 3275
 3280
 3285
 3290
 3295
 3300
 3305
 3310
 3315
 3320
 3325
 3330
 3335
 3340
 3345
 3350
 3355
 3360
 3365
 3370
 3375
 3380
 3385
 3390
 3395
 3400
 3405
 3410
 3415
 3420
 3425
 3430
 3435
 3440
 3445
 3450
 3455
 3460
 3465
 3470
 3475
 3480
 3485
 3490
 3495
 3500
 3505
 3510
 3515
 3520
 3525
 3530
 3535
 3540
 3545
 3550
 3555
 3560
 3565
 3570
 3575
 3580
 3585
 3590
 3595
 3600
 3605
 3610
 3615
 3620
 3625
 3630
 3635
 3640
 3645
 3650
 3655
 3660
 3665
 3670
 3675
 3680
 3685
 3690
 3695
 3700
 3705
 3710
 3715
 3720
 3725
 3730
 3735
 3740
 3745
 3750
 3755
 3760
 3765
 3770
 3775
 3780
 3785
 3790
 3795
 3800
 3805
 3810
 3815
 3820
 3825
 3830
 3835
 3840
 3845
 3850
 3855
 3860
 3865
 3870
 3875
 3880
 3885
 3890
 3895
 3900
 3905
 3910
 3915
 3920
 3925
 3930
 3935
 3940
 3945
 3950
 3955
 3960
 3965
 3970
 3975
 3980
 3985
 3990
 3995
 4000
 4005
 4010
 4015
 4020
 4025
 4030
 4035
 4040
 4045
 4050
 4055
 4060
 4065
 4070
 4075
 4080
 4085
 4090
 4095
 4100
 4105
 4110
 4115
 4120
 4125
 4130
 4135
 4140
 4145
 4150
 4155
 4160
 4165
 4170
 4175
 4180
 4185
 4190
 4195
 4200
 4205
 4210
 4215
 4220
 4225
 4230
 4235
 4240
 4245
 4250
 4255
 4260
 4265
 4270
 4275
 4280
 4285
 4290
 4295
 4300
 4305
 4310
 4315
 4320
 4325
 4330
 4335
 4340
 4345
 4350
 4355
 4360
 4365
 4370
 4375
 4380
 4385
 4390
 4395
 4400
 4405
 4410
 4415
 4420
 4425
 4430
 4435
 4440
 4445
 4450
 4455
 4460
 4465
 4470
 4475
 4480
 4485
 4490
 4495
 4500
 4505
 4510
 4515
 4520
 4525
 4530
 4535
 4540
 4545
 4550
 4555
 4560
 4565
 4570
 4575
 4580
 4585
 4590
 4595
 4600
 4605
 4610
 4615
 4620
 4625
 4630
 4635
 4640
 4645
 4650
 4655
 4660
 4665
 4670
 4675
 4680
 4685
 4690
 4695
 4700
 4705
 4710
 4715
 4720
 4725
 4730
 4735
 4740
 4745
 4750
 4755
 4760
 4765
 4770
 4775
 4780
 4785
 4790
 4795
 4800
 4805
 4810
 4815
 4820
 4825
 4830
 4835
 4840
 4845
 4850
 4855
 4860
 4865
 4870
 4875
 4880
 4885
 4890
 4895
 4900
 4905
 4910
 4915
 4920
 4925
 4930
 4935
 4940
 4945
 4950
 4955
 4960
 4965
 4970
 4975
 4980
 4985
 4990
 4995
 5000
 5005
 5010
 5015
 5020
 5025
 5030
 5035
 5040
 5045
 5050
 5055
 5060
 5065
 5070
 5075
 5080
 5085
 5090
 5095
 5100
 5105
 5110
 5115
 5120
 5125
 5130
 5135
 5140
 5145
 5150
 5155
 5160
 5165
 5170
 5175
 5180
 5185
 5190
 5195
 5200
 5205
 5210
 5215
 5220
 5225
 5230
 5235
 5240
 5245
 5250
 5255
 5260
 5265
 5270
 5275
 5280
 5285
 5290
 5295
 5300
 5305
 5310
 5315
 5320
 5325
 5330
 5335
 5340
 5345
 5350
 5355
 5360
 5365
 5370
 5375
 5380
 5385
 5390
 5395
 5400
 5405
 5410
 5415
 5420
 5425
 5430
 5435
 5440
 5445
 5450
 5455
 5460
 5465
 5470
 5475
 5480
 5485
 5490
 5495
 5500
 5505
 5510
 5515
 5520
 5525
 5530
 5535
 5540
 5545
 5550
 5555
 5560
 5565
 5570
 5575
 5580
 5585
 5590
 5595
 5600
 5605
 5610
 5615
 5620
 5625
 5630
 5635
 5640
 5645
 5650
 5655
 5660
 5665
 5670
 5675
 5680
 5685
 5690
 5695
 5700
 5705
 5710
 5715
 5720
 5725
 5730
 5735
 5740
 5745
 5750
 5755
 5760
 5765
 5770
 5775
 5780
 5785
 5790
 5795
 5800
 5805
 5810
 5815
 5820
 5825
 5830
 5835
 5840
 5845
 5850
 5855
 5860
 5865
 5870
 5875
 5880
 5885
 5890
 5895
 5900
 5905
 5910
 5915
 5920
 5925
 5930
 5935
 5940
 5945
 5950
 5955
 5960
 5965
 5970
 5975
 5980
 5985
 5990
 5995
 6000
 6005
 6010
 6015
 6020
 6025
 6030
 6035
 6040
 6045
 6050
 6055
 6060
 6065
 6070
 6075
 6080
 6085
 6090
 6095
 6100
 6105
 6110
 6115
 6120
 6125
 6130
 6135
 6140
 6145
 6150
 6155
 6160
 6165
 6170
 6175
 6180
 6185
 6190
 6195
 6200
 6205
 6210
 6215
 6220
 6225
 6230
 6235
 6240
 6245
 6250
 6255
 6260
 6265
 6270
 6275
 6280
 6285
 6290
 6295
 6300
 6305
 6310
 6315
 6320
 6325
 6330
 6335
 6340
 6345
 6350
 6355
 6360
 6365
 6370
 6375
 6380
 6385
 6390
 6395
 6400
 6405
 6410
 6415
 6420
 6425
 6430
 6435
 6440
 6445
 6450
 6455
 6460
 6465
 6470
 6475
 6480
 6485
 6490
 6495
 6500
 6505
 6510
 6515
 6520
 6525
 653


used, these builders comprise from about 5% to about 80% by weight of the detergent compositions.

Inorganic detergent builders include, but are not limited to, the alkali metal, ammonium and alkanolammonium salts of polyphosphates (exemplified by the tripolyphosphates, pyrophosphates, and glassy polymeric meta-phosphates), phosphonates, phytic acid, silicates, carbonates (including bicarbonates and sesquicarbonates), sulphates, and aluminosilicates. However, non-phosphate builders are required in some locales.

Examples of silicate builders are the alkali metal silicates, particularly those having a $\text{SiO}_2:\text{Na}_2\text{O}$ ratio in the range 1.6:1 to 3.2:1 and layered silicates, such as the layered sodium silicates described in U.S. Patent 4,664,839, issued May 12, 1987 to H. P. Rieck, available from Hoechst under the trademark "SKS"; SKS-6 is an especially preferred layered silicate builder.

Carbonate builders, especially a finely ground calcium carbonate with surface area greater than $10 \text{ m}^2/\text{g}$, are preferred builders that can be used in granular compositions. The density of such alkali metal carbonate built detergents can be in the range of 450-850 g/l with the moisture content preferably below 4%. Examples of carbonate builders are the alkaline earth and alkali metal carbonates as disclosed in German Patent Application No. 2,321,001 published on November 15, 1973.

Aluminosilicate builders are especially useful in the present invention. Preferred aluminosilicates are zeolite builders which have the formula:



wherein z and y are integers of at least 6, the molar ratio of z to y is in the range from 1.0 to about 0.5, and x is an integer from about 15 to about 264.

Useful aluminosilicate ion exchange materials are commercially available. These aluminosilicates can be crystalline or amorphous in structure and can be naturally-occurring aluminosilicates or synthetically derived. A method for producing aluminosilicate ion exchange materials is disclosed in U.S. Patent 3,985,669, Krummel, et al, issued October 12, 1976. Preferred synthetic crystalline aluminosilicate ion exchange materials useful herein are available under the designations Zeolite A, Zeolite P (B), and Zeolite X. Preferably, the aluminosilicate has a particle size of about 0.1-10 microns in diameter.

Organic detergent builders suitable for the purposes of the present invention include, but are not restricted to, a wide variety of polycarboxylate compounds, such as ether polycarboxylates, including oxydisuccinate, as disclosed in Berg, U.S. Patent 3,128,287, issued April 7, 1964, and Lamberti et al, U.S. Patent 3,635,830, issued January 18, 1972. See also "TMS/TDS" builders of U.S. Patent 4,663,071, issued to Bush et al, on May 5, 1987.

Other useful detergent builders include the ether hydroxypolycarboxylates, copolymers of maleic anhydride with ethylene or vinyl methyl ether, 1, 3, 5-trihydroxy benzene-2, 4, 6-trisulphonic acid, and carboxymethyloxysuccinic acid, the various alkali metal, ammonium and substituted ammonium salts of polyacetic acids such as ethylenediamine tetraacetic acid and nitrilotriacetic acid, as well as polycarboxylates such as mellitic acid, succinic acid, oxydisuccinic acid, polymaleic acid, benzene 1,3,5-tricarboxylic acid, carboxymethyloxysuccinic acid, and soluble salts thereof.

Citrate builders, e.g., citric acid and soluble salts thereof (particularly sodium salt), are preferred polycarboxylate builders that can also be used in granular compositions, especially in combination with zeolite and/or layered silicate builders.

Also suitable in the detergent compositions of the present invention are the 3,3-dicarboxy-4-oxa-1,6-hexanedioates and the related compounds disclosed in U.S. Patent 4,566,984, Bush, issued January 28, 1986.

In situations where phosphorus-based builders can be used, and especially in the formulation of bars used for hand-laundrying operations, the various alkali metal phosphates such as the well-known sodium tripolyphosphates, sodium pyrophosphate and sodium orthophosphate can be used. Phosphonate builders such as ethane-1-hydroxy-1,1-diphosphonate and other known phosphonates (see, for example, U.S. Patents 3,159,581; 3,213,030; 3,422,021; 3,400,148 and 3,422,137) can also be used.

Optional Detergent Adjuncts

As a preferred embodiment, the conventional detergent ingredients employed herein can be selected from typical detergent composition components such as detergent surfactants and detergent builders. Optionally, the detergent ingredients can include one or more other detergent adjuncts or other materials for assisting or enhancing cleaning performance, treatment of the substrate to be cleaned, or to modify the aesthetics of the detergent composition. Usual detergent adjuncts of detergent compositions include the ingredients set forth in U.S. Pat. No. 3,936,537, Baskerville et al. Adjuncts which can also be included in detergent compositions employed in the present invention, in their conventional art-established levels for use (generally from 0% to about 20% of the detergent ingredients, preferably from about 0.5% to about 10%), include enzymes, especially proteases, lipases and cellulases, color speckles, suds boosters, suds suppressors, antitarnish and/or anticorrosion agents, soil-suspending agents, soil release agents, dyes, fillers, optical brighteners, germicides, alkalinity sources, hydrotropes, antioxidants, enzyme stabilizing agents, perfumes, solvents, solubilizing agents, clay soil removal/anti-redeposition agents, polymeric dispersing agents, processing aids, fabric softening components such as smectite clays, static control agents, etc.

Bleach systems optionally, but preferably, will also comprise a chelant which not only enhances bleach stability by scavenging heavy metal ions which tend to decompose bleaches, but also assists in the removal of polyphenolic stains such as tea stains, and the like. Various chelants, including the aminophosphonates, available as DEQUEST
5 from Monsanto, the nitrilotriacetates, the hydroxyethyl-ethylenediamine triacetates, and the like, are known for such use. Preferred biodegradable, non-phosphorus chelants include ethylenediamine disuccinate ("EDDS"; see U.S. Patent 4,704,233, Hartman and Perkins), ethylenediamine-N,N'-diglutamate (EDDG) and 2-hydroxypropylenediamine-N,N'-disuccinate (HPDDS) compounds. Such chelants can be used in their alkali or
10 alkaline earth metal salts, typically at levels from about 0.1% to about 10% of the present compositions.

Optionally, the detergent compositions employed herein can comprise, in addition to the bleaching system of the present invention, one or more other conventional bleaching agents, activators, or stabilizers. In general, the formulator will ensure that
15 the bleach compounds used are compatible with the detergent formulation. Conventional tests, such as tests of bleach activity on storage in the presence of the separate or fully-formulated ingredients, can be used for this purpose.

Specific examples of optional bleach activators for incorporation in this invention include tetraacetyl ethylene diamine (TAED), N-acyl caprolactams, alkanoyloxy-
20 benzenesulfonates, including nonanoyloxybenzenesulfonate and benzoyloxybenzenesulfonate, the benzoxazin-type bleaching activators disclosed in U.S. Patent 4,966,723, Hodge et al, issued Oct. 30, 1990, and the peroxyacid agents and activators having amide moieties disclosed in U.S. Patent 4,634,551, Burns et al, issued Jan. 6, 1987. Such bleaching compounds and agents can be optionally included in detergent
25 compositions in their conventional art-established levels of use, generally from 0% to about 15%, by weight of detergent composition.

Bleaching activators of the invention are especially useful in conventional laundry detergent compositions such as those typically found in granular detergents or laundry bars. U.S. Patent 3,178,370, Okenfuss, issued April 13, 1965, describes laundry
30 detergent bars and processes for making them. Philippine Patent 13,778, Anderson, issued Sept. 23, 1980, describes synthetic detergent laundry bars. Methods for making laundry detergent bars by various extrusion methods are well known in the art.

Perhydrolysis Speed and Yield

The activators herein have better perhydrolysis speed and yield when compared
35 to other lactam derived bleach activators. For example, after approximately five minutes, a solution of nonanoyl valerolactam generates approximately 95% of the theoretical yield of peracid. In comparison, after approximately five minutes, an identical solution of nonanoyl caprolactam generates approximately 35% of the


theoretical yield of peracid. After 15 minutes, the nonanoyl caprolactam solution generates approximately 68% of the theoretical yield of peracid. Accordingly, the user can expect better bleaching performance at shorter wash times with the valerolactam activators.

- 5 The comparison is made using standard peracid titration procedures. The activator is dispersed in a detergent solution. At specific time intervals, samples of the solution are taken and added to a mixture of acetic acid, water, and ice to quench the perhydrolysis reaction. A quantity of potassium iodide is added to the sample giving the sample a brown color. The resulting solution is titrated with thiosulphate until the
10 color is removed. From the known quantity of thiosulphate used in the titration, the amount of peracid can be determined.


Liquid Bleaches

The bleach activators of this invention are also useful in liquid bleach compositions. Therefore, in accordance with one aspect of the invention, a stable
15 aqueous liquid bleach is provided. In a preferred embodiment, the liquid bleaches comprise a liquid valerolactam such as decanoyl valerolactam or nonanoyl valerolactam which is most preferably emulsified in the peroxide liquid composition. Such liquid bleaches comprise:

- A) from about 1% to about 25%, more preferably from about 3% to about 12%,
20 most preferably from about 5% to about 10%, by weight, of a (solid or, preferably, liquid) valerolactam bleach activator selected from the group consisting of:



wherein R_1 is as defined above;



wherein R^2 , R^3 , R^4 , R^5 , and R^6 are as defined above; and

- 35 c) mixtures of a) and b);
- B) from about 0.1% to about 10%, more preferably from about 0.3% to about 7%, most preferably from about 0.5% to about 5%, by weight, of a peroxygen bleaching compound comprising hydrogen peroxide or which is capable of yielding hydrogen peroxide in an aqueous solution.

Optionally, the liquid bleach composition can further comprise from about 1% to about 20%, preferably from about 5% to about 15%, by weight, of a phase stabilizer and from about 0.001% to about 2%, preferably from about 0.05% to about 1%, by weight, of a chelating agent. The balance of the liquid bleach composition is water.

5 The liquid peroxide-containing bleach is formulated in the acid pH range for stability. In-use in a laundering operation, the bleach is added to a laundry liquor which typically has a pH in the base range, i.e., 9.5 - 13, which then destabilizes the peroxide to perform its bleaching function.

Nonionic Surfactant

10 The liquid bleach compositions of the invention, optionally, but preferably, include a nonionic surfactant as a phase stabilizer to facilitate maintenance of its continuous isotropic state. To this end, several nonionic surfactants are particularly useful. Suitable nonionic surfactants include the polyethylene oxide condensates of alkyl phenols, e.g., the condensation products of alkyl phenols having an alkyl group
15 containing from about 6 to 15 carbon atoms, in either a straight chain or branched chain configuration, with from about 3 to 20 moles of ethylene oxide per mole of alkyl phenol.

Other nonionic surfactants which function as suitable phase stabilizers are the water-soluble and water-dispersible condensation products of aliphatic alcohols
20 containing from 8 to 22 carbon atoms, in either straight chain or branched configuration, with from 3 to 20 moles of ethylene oxide per mole of alcohol. Still other nonionic surfactants include semi-polar nonionic surfactants such as water-soluble amine oxides containing one alkyl moiety of from about 10 to 18 carbon atoms and two moieties selected from about 1 to about 3 carbon atoms; water-soluble phosphine
25 oxides containing one alkyl moiety of about 10 to 18 carbon atoms and two moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to 3 carbon atoms; and water-soluble sulfoxides containing only alkyl moiety of from about 10 to 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to 3 carbon atoms.

30 Preferred nonionic surfactants are of the formula $R^1(OC_2H_4)_nOH$, wherein R^1 is a C_8 - C_{16} alkyl group or C_8 - C_{12} alkyl phenyl group, and n is from 3 to about 20. Particularly preferred are condensation products of C_9 - C_{15} alcohols with from about 5 to about 20 moles of ethylene oxide per mole of alcohol. The most preferred nonionic of this type is an alkyl ethoxylate having from about 9 to 11 carbon atoms and an
35 average degree of ethoxylation of about 10 which is available from Shell Oil Co. under the product name of NEODOL 91-10.

The liquid bleach composition will generally comprise from about 1% to about 20%, preferably from about 5% to about 15%, by weight, of the nonionic surfactant phase stabilizer.

With the aforementioned chelating agent and phase stabilizer, i.e. nonionic surfactant, the stable aqueous liquid bleach composition in accordance with the invention can be produced. The resulting liquid bleach composition has a relatively low viscosity which renders it more pourable and therefore, more convenient for users especially when the composition is used as an additive. The viscosity of the present liquid bleach is preferably in a range from about 10 to about 500 cps, more preferably from about 10 to about 300 cps, and most preferably from about 10 to 100 cps.

pH Adjusting Agent

It has been found that optimum stability of the liquid peroxygen bleaches is achieved when the aqueous liquid bleach has a pH in range from about 2 to about 7, more preferably from about 3 to about 5, and most preferably from about 3.5 to about 4.5. For purposes of achieving such pH's in the present stable aqueous liquid bleach composition, a pH adjusting agent may optionally be included. It is a well known technique to use pH adjusting agents to alter aqueous solutions such as the present liquid bleach, to the desired pH.

Typical pH adjusting agents can be either of the acid type or of the base type. Acidic pH adjusting agents are designed to compensate for the presence of other highly alkaline materials and include organic and inorganic acids, acid mixtures and acid salts. Non-limiting examples of such acidic pH adjusting agents include citric acid, glycolic acid, phosphoric acid, lauric acid and mixtures thereof. Representative examples of alkaline pH adjusting agents include but not limited to sodium hydroxide, salts of phosphates, citrates and mixtures thereof.

In addition to the materials described above, the liquid bleach may also include perfumes colorants, brighteners, viscosity adjusters such as thickeners, and other conventional components typically used in detergent compositions, if compatible.

The liquid bleach of the invention can be produced by a wide variety of processes. While not intending to be limiting, the most economical and easiest manner in which the liquid bleach can be produced is to simply disperse all of the preferred components in water. As those skilled in the art will appreciate, it may be desirable to disperse certain components in water before others. This offers an inexpensive way to produce the present liquid bleach composition.

In accordance with another aspect of the invention, a method of bleaching fabrics comprises the step of contacting fabrics with a diluted aqueous solution of the liquid composition of the invention. Another method contemplated by the invention involves laundering soiled clothes, using the liquid bleach composition as an additive. The

method comprises the steps of contacting fabrics with an effective amount of a detergent composition in combination with an effective amount of a stable aqueous liquid bleach composition. In practicing these methods, the stable aqueous liquid bleach compositions of the present invention can be used in widely varying concentrations depending on the particular application involved but are generally utilized in an amount sufficient to provide from about 1.0 ppm to about 50 ppm available oxygen from the peracid in solution.

The following examples are given to further illustrate the present invention, but are not intended to be limiting thereof.

EXAMPLE I

Synthesis of benzoyl valerolactam - To a 500 ml flask equipped with a reflux condenser, overhead stirrer, and addition funnel is charged 0.12 moles of valerolactam, 0.15 moles of triethylamine and 150 ml of toluene. The solution is heated to 80°C and a solution of 0.12 moles of benzoyl chloride dissolved in 50 ml of toluene is added over 20 minutes. The mixture is refluxed for six hours with stirring, cooled, and filtered. The filtrate is concentrated under vacuum to a brown solid which is recrystallized from toluene to yield 0.092 moles of benzoyl valerolactam. ¹H NMR analysis shows the product to be greater than 95% pure. Analysis: ¹H NMR (ppm) - 1.9(M, 4H); 2.5(t, 2H); 3.7(t, 2H); 7.4(M, 5H). IR (cm⁻¹) - 2925, 2856, 1686, 1673, 1458, 1287.

EXAMPLE II

Synthesis of nonanoyl valerolactam - To a 5 liter flask equipped with a reflux condenser, argon sweep, addition funnel, and overhead stirrer is charged 2.5 moles of valerolactam, 2.75 moles of triethylamine, and 2 liters of toluene. The solution is heated to reflux, and 2.5 moles of nonanoyl chloride is added over 1.5 hours. The mixture is refluxed with stirring for 6 hours, cooled, and filtered. The filtrate is concentrated under vacuum, and the resultant oil is distilled at 162°C and approximately 1 mm Hg to yield 1.6 moles of nonanoyl valerolactam. NMR analysis shows the product to be greater than 95% pure. Analysis: ¹H NMR (ppm) - 0.9(t, 3H); 1.3(M, 10H); 1.6(M, 2H); 1.9(M, 4H); 2.6(M, 2H); 2.9(t, 2H); 3.7(M, 2H). IR (cm⁻¹) - 2927, 2857, 1697, 1461, 1377, 1291, 1197, 1161.

EXAMPLE III

A granular detergent composition is prepared comprising the following ingredients.

	<u>Component</u>	<u>Weight %</u>
35	C ₁₂ linear alkyl benzene sulfonate	22
	Phosphate (as sodium tripolyphosphate)	20
	Sodium carbonate	10
	Sodium silicate	3

	Sodium percarbonate*	20
	Ethylenediamine disuccinate chelant (EDDS)	0.4
	Sodium sulfate	5.5
	Nonanoyl valerolactam	10
5	Minors, filler** and water	Balance to 100%

* Average particle size of 400 to 1200 microns.

**Can be selected from convenient materials such as CaCO_3 , talc, clay, silicates, and the like.

10 An aqueous crutcher mix of heat and alkali stable components of the detergent composition is prepared and spray-dried. The other ingredients are admixed therewith so that they contain the ingredients tabulated at the levels shown.

The detergent granules with bleaching system are added together with a 6 lb. (2.7 kg) load of fabrics to a Sears KENMORE washing machine. Actual weight of the detergent composition is taken to provide a 1500 ppm concentration of the detergent
15 composition in the 17 gallon (65 l) water-fill machine. The water used has 7 grains/gallon hardness and a pH of 7 to 7.5 prior to (about 9 to about 10.5 after) addition of the detergent composition. The fabrics are laundered at 35°C (95°F) for a full cycle (12 min.) and rinsed at 21°C (70°F).

At the end of the last rinse cycle, the test swatches are dried in a dryer.
20 Tristimulus meter readings (L,a,b) are then determined for each test swatch. Whiteness performance in terms of Hunter Whiteness Values (W) is then calculated according to the following equation:

$$W = (7L^2 - 40Lb)/700$$

25 The higher the value for W, the better the whiteness performance. In the above test, fabrics exposed to the bleaching system display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE IV

30 A granular detergent is prepared by a procedure identical to that of Example III, with the exception that 20% benzoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator. The laundering method of Example III is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE V

5 A granular detergent is prepared by a procedure identical to that of Example III, with the exceptions that 10% 4-nitrobenzoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator. The laundering method of Example III is repeated. In the test, all fabrics display significantly improved whiteness after

laundrying compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE VI

5 A granular detergent is prepared by a procedure identical to that of Example III, with the exceptions that 5% isononanoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator and the amount of sodium percarbonate is 10%. The laundrying method of Example III is repeated. In the test, all fabrics display significantly improved whiteness after laundrying compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE VII

10 A granular detergent is prepared by a procedure identical to that of Example III, with the exceptions that 20% 3-chlorobenzoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator and the amount of sodium percarbonate is 20%. The laundrying method of Example III is repeated. In the test, all fabrics display significantly improved whiteness after laundrying compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE VIII

20 A granular detergent is prepared by a procedure identical to that of Example III, with the single exception that 15% of a 1:1 mixture of benzoyl valerolactam and nonanoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator. The laundrying method of Example III is repeated. In the test, all fabrics display significantly improved whiteness after laundrying compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE IX

25 A granular detergent is prepared by a procedure identical to that of Example III, with the single exception that 15% of a 1:1 mixture of benzoyl valerolactam and nonanoyloxybenzenesulfonate is substituted for the nonanoyl valerolactam bleach activator. The laundrying method of Example III is repeated. In the test, all fabrics display significantly improved whiteness after laundrying compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE X

35 A granular detergent is prepared by a procedure identical to that of Example III, with the single exception that 15% of a 1:1 mixture of 4-ethylbenzoyl valerolactam and a benzoxazin-type bleach activator is substituted for the nonanoyl valerolactam bleach activator. The laundrying method of Example III is repeated. In the test, all fabrics display significantly improved whiteness after laundrying compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XI

A granular detergent is prepared by a procedure identical to that of Example III, with the single exception that an equivalent amount of decanoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator. The laundering method of
 5 Example III is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XII

A granular detergent composition is prepared comprising the following
 10 ingredients.

	<u>Component</u>	<u>Weight %</u>
	Anionic alkyl sulfate	7
	Nonionic surfactant	5
	Zeolite A (0.1-10 micron)	10
15	Trisodium citrate	2
	SKS-6 silicate builder	10
	Acrylate maleate polymer	4
	Nonanoyl valerolactam	10
	Sodium percarbonate	25
20	Sodium carbonate	5
	Ethylenediamine disuccinate chelant (EDDS)	0.4
	Suds suppressor	2
	Enzymes*	1.5
	Soil release agent	0.2
25	Minors, filler** and water	Balance to 100%

*1:1:1 mixture of protease, lipase, and cellulase.

**Can be selected from convenient materials such as CaCO_3 , talc, clay, silicates, and the like.

An aqueous crutcher mix of heat and alkali stable components of the detergent
 30 composition is prepared and spray-dried. The other ingredients are admixed therewith so that they contain the ingredients tabulated at the levels shown.

The detergent granules with bleaching system are added together with a 2.7 kg load of fabrics to an automatic washing machine. Actual weights of detergent and ester compositions are taken to provide a 5000 ppm concentration of the detergent
 35 composition in the 17 liter (4.5 gallon) water-fill machine. The water used has 10 grains/ gallon hardness and a pH of 7 to 7.5 prior to (about 9 to about 10.5 after) addition of the detergent composition.

The fabrics are laundered at 40°C (104°F) for a full cycle (40 min.) and rinsed at 21°C (70°F).

At the end of the last rinse cycle, the test swatches are dried in a dryer. Tristimulus meter readings (L,a,b) are then determined for each test swatch. Whiteness performance in terms of Hunter Whiteness Values (W) is then calculated according to the following equation:

$$W = (7L^2 - 40Lb)/700$$

The higher the value for W, the better the whiteness performance. In the above test, fabrics exposed to the bleaching system display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XIII

A granular detergent is prepared by a procedure identical to that of Example XII, with the exception that 10% isononanoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator. The laundering method of Example XII is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XIV

A granular detergent is prepared by a procedure identical to that of Example XII, with the exceptions that 10% 2-ethylhexanoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator. The laundering method of Example XII is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XV

A granular detergent is prepared by a procedure identical to that of Example XII, with the exceptions that 15% benzoyl valerolactam is substituted for the nonanoyl valerolactam bleach activator and the amount of sodium percarbonate is 30%. The laundering method of Example XII is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XVI

A granular detergent is prepared by a procedure identical to that of Example XII, with the single exception that 15% of a 1:1 mixture of 4-chlorobenzoyl valerolactam and a benzoxazin-type bleach activator is substituted for the nonanoyl valerolactam bleach activator. The laundering method of Example XII is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XVII

A granular detergent is prepared by a procedure identical to that of Example XII, with the single exception that 15% of a 1:1 mixture of isononanoyl valerolactam and tetraacetyl ethylene diamine bleach activator is substituted for the nonanoyl
 5 valerolactam bleach activator. The laundering method of Example XII is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XVIII

A granular detergent is prepared by a procedure identical to that of Example XII,
 10 with the single exception that 15% of a 1:1 mixture of isononanoyl valerolactam and benzoyl caprolactam is substituted for the nonanoyl valerolactam bleach activator. The laundering method of Example XII is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XIX

15 A laundry bar suitable for hand-washing soiled fabrics is prepared comprising the following ingredients.

	<u>Component</u>	<u>Weight %</u>
	C ₁₂ linear alkyl benzene sulfonate	30
20	Phosphate (as sodium tripolyphosphate)	7
	Sodium carbonate	15
	Sodium pyrophosphate	7
	Coconut monoethanolamide	2
	Zeolite A (0.1-10 microns)	5
25	Carboxymethylcellulose	0.2
	Polyacrylate (m.w. 1400)	0.2
	Benzoyl valerolactam	6.5
	Sodium percarbonate	15
	Brightener, perfume	0.2
30	Protease	0.3
	CaSO ₄	1
	MgSO ₄	1
	Water and Filler*	Balance to 100%

*Can be selected from convenient materials such as CaCO₃, talc, clay, silicates, and the
 35 like.

Detergent laundry bar is extruded in conventional soap or detergent bar making equipment as commonly used in the art. Testing is conducted following the methods used in Example IV. In the test, fabrics exposed to the bleaching system display

significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XX

5 A laundry bar is prepared by a procedure identical to that of Example XIX, with the two exceptions that 15% of a 1:1 mixture of benzoyl valerolactam and nonanoyl valerolactam is substituted for the benzoyl valerolactam bleach activator, and the level of sodium percarbonate is increased to 20%. The laundering method of Example XIX is repeated. In the test, all fabrics display significantly improved whiteness after
10 laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XXI

A laundry bar is prepared by a procedure identical to that of Example XIX, with the two exceptions that 15% of a 1:1 mixture of isononanoyl valerolactam and 2,4-dichlorobenzoyl valerolactam is substituted for the benzoyl valerolactam bleach
15 activator, and the level of sodium percarbonate is increased to 20%. The laundering method of Example XIX is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XXII

20 A laundry bar is prepared by a procedure identical to that of Example XIX, with the two exceptions that 15% of a 1:1 mixture of decanoyl valerolactam and benzoyl valerolactam is substituted for the benzoyl valerolactam bleach activator, and the level of sodium percarbonate is increased to 20%. The laundering method of Example XIX is repeated. In the test, all fabrics display significantly improved whiteness after
25 laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XXIII

A laundry bar is prepared by a procedure identical to that of Example XIX, with the two exceptions that an equivalent amount of butoxybenzoyl valerolactam is
30 substituted for the benzoyl valero- lactam bleach activator, and the level of sodium percarbonate is increased to 20%. The laundering method of Example XIX is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XXIV

35 A bleaching system is prepared comprising the following ingredients.

<u>Component</u>	<u>Weight %</u>
Nonanoyl valerolactam	15

Tetraacetyl ethylene diamine	15
Sodium percarbonate	45
Chelant (ethylenediamine disuccinate, EDDS)	10
Filler* and water	Balance to 100%

- 5 *Can be selected from convenient materials such as CaCO_3 , talc, clay, silicates, and the like.

Testing is conducted following the methods used in Example IV with the single exception that the an equivalent amount of the above bleaching system is substituted for the detergent composition used in Example IV. In the test, fabrics exposed to the
10 bleaching system display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XXV

A bleaching system is prepared by a procedure identical to that of Example XXIV, with the exception that an equivalent amount of 4-butylbenzoyl valerolactam is
15 substituted for the nonanoyl valerolactam bleach activator. The laundering method of Example XXIV is repeated. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XXVI

20 A bleaching system is prepared by a procedure identical to that of Example XXIV, with the exception that an equivalent amount of terephthaloyl divalerylactam is substituted for the nonanoyl valerolactam bleach activator. The laundering method of Example XXIV is repeated. In the test, all fabrics display significantly improved
25 whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

While the compositions and processes of the present invention are useful in conventional laundering operations, it is to be understood that they are also useful in any cleaning system which involves low water: fabric ratios. One such system is disclosed in U.S. Patent 4,489,455, Spindel, issued Dec. 25, 1984, which involves a
30 washing machine apparatus which contacts fabrics with wash water containing detergent ingredients using a low water: fabric ratio rather than the conventional method of immersing fabrics in an aqueous bath. The compositions herein provide excellent bleaching performance in such mechanical systems. Typically, the ratio of water: fabric ranges from about 0.5:1 to about 6:1 (liters of water: kg of fabric).

EXAMPLE XXVII

35 Using the machine and operating conditions disclosed in U.S. Patent 4,489,455, cited above, 25 grams of a composition according to Example IV herein are used to launder fabrics with concurrent bleaching. If desired, sudsing of the composition can be

minimized by incorporating therein from 0.2% to 2% by weight of a fatty acid, secondary alcohol, or silicone suds controlling ingredient. In the test, all fabrics display significantly improved whiteness after laundering compared with fabrics which have not been exposed to the bleaching system of the invention.

- 5 The valerolactam bleach activators are preferably not absorbed onto the peroxygen bleaching compound. To do so in the presence of other organic detergent ingredients could cause safety problems. It has now been discovered that the acyl valerolactam bleach activators of this invention can be dry-mixed with peroxygen bleaching compounds, especially perborate and percarbonate, and thereby avoid
10 potential safety problems. If the valerolactam is a liquid, before dry-mixing, it can be formed into a suitable "particle" using conventional means known in the art.

EXAMPLE XXVIII

A laundry bar suitable for hand-washing soiled fabrics is prepared comprising the following ingredients.

	<u>Component</u>	<u>Weight %</u>
15	Linear alkyl benzene sulfonate	30
	Phosphate (as sodium tripolyphosphate)	7
	Sodium carbonate	20
	Sodium pyrophosphate	7
20	Coconut monoethanolamide	2
	Zeolite A (0.1-10 microns)	5
	Carboxymethylcellulose	0.2
	Polyacrylate (m.w. 1400)	0.2
	Isononanoyl valerolactam	5
25	Sodium perborate tetrahydrate	10
	Brightener, perfume	0.2
	Protease	0.3
	CaSO ₄	1
	MgSO ₄	1
30	Water	4
	Filler*	Balance to 100%

*Can be selected from convenient materials such as CaCO₃, talc, clay, silicates, and the like.

- 35 A detergent laundry bar is extruded in conventional soap or detergent bar making equipment as commonly used in the art with the bleaching activator dry-mixed with the perborate bleaching compound and not affixed to the surface of the perborate. Testing is conducted following the methods used in Example IV. In the test, fabrics exposed to the bleaching system display significantly improved whiteness after laundering

compared with fabrics which have not been exposed to the bleaching system of the invention.

EXAMPLE XXIX

A granular detergent composition is prepared comprising the following
5 ingredients.

	<u>Component</u>	<u>Weight %</u>
	Linear alkyl benzene sulfonate	20
	Phosphate (as sodium tripolyphosphate)	20
	Sodium carbonate	10
10	Sodium silicate	3
	Sodium perborate tetrahydrate	20
	Ethylenediamine disuccinate chelant (EDDS)	0.4
	Sodium sulfate	5.5
	4-nitrobenzoyl valerolactam	5
15	Nonanoyloxybenzenesulfonate	5
	Minors, filler** and water	Balance to 100%

**Can be selected from convenient materials such as CaCO_3 , talc, clay, silicates, and the like.

Aqueous crutcher mixes of heat and alkali stable components of the detergent
20 compositions are prepared and spray-dried and the other ingredients are dry-mixed so that they contain the ingredients tabulated at the levels shown.

Testing is conducted following the methods used in Example V. In the test,
fabrics exposed to the bleaching system display significantly improved whiteness after
laundering compared with fabrics which have not been exposed to the bleaching system
25 of the invention.

EXAMPLE XXX

This Example illustrates several liquid bleach compositions in accordance with
the invention, all of which are made by the general process described hereinafter. The
desired amount of a chelating agent is added to a beaker of water, after which the
30 resulting solution is stirred until the chelating agent is completely dissolved. A phase
stabilizer is added to the solution while it is being continuously stirred. Thereafter, the
bleach activator and optionally an additional chelating agent is dissolved in the solution.
The pH of the solution is adjusted to about 4.0 with an alkaline adjusting agent such as
sodium hydroxide.

35 The following translucent, stable aqueous liquid bleach compositions (Samples
A-F) are made as described above, all amounts being expressed as percentages by
weight.

TABLE I

Samples

	A	B	C	D	E	F
Water	81.28	81.86	82.44	83.02	78.60	83.98
NEODOL 91-10 ¹	10.00	10.00	10.00	10.00	10.00	10.00
Dipicolinic Acid ²	0.05	0.05	0.05	0.05	0.05	0.05
DEQUEST 2010 ³	0.05	0.05	0.05	0.05	0.05	0.05
Nonanoyl Valerolactam	5.80	5.80	5.80	5.80	7.71	3.87
Citric Acid	0.50	0.50	0.50	0.50	0.50	0.50
NaOH	to pH4	to pH4	to pH4	to pH4	to pH4	to pH4
Hydrogen Peroxide	2.32	1.74	1.16	0.58	3.09	1.55


5 ¹ Alkyl ethoxylate available from The Shell Oil Company.

² 2,6-Pyridine dicarboxylic acid commercially available from Aldrich Chemical Co.


³ Hydroxy-ethylidene diphosphonic acid commercially available from Monsanto Co.

What is Claimed is:


1. An acyl valerolactam bleach activator selected from the group consisting of:



wherein R^1 is a substituted or unsubstituted alkyl or alkoxy group containing from about 1 to about 18 carbon atoms wherein the longest linear alkyl or alkoxy chain extending from and including the carbonyl carbon contains from about 2 to about 12 carbon atoms;



wherein R^2 , R^3 , R^4 , R^5 , and R^6 are members selected from the group consisting of H, halogen, alkyl, alkoxy, alkoxyaryl, alkaryl, and alkaryloxy moieties containing from about 1 to about 12 carbon atoms, and substituents having the structures:



wherein R^7 is selected from the group consisting of H, alkyl, alkaryl, alkoxy, alkoxyaryl, alkaryloxy, and aminoalkyl; X is O, NH, or NR^9 , wherein R^9 is H or a C_1 - C_4 alkyl group; and R^8 is an alkyl, cycloalkyl, or aryl group containing from 3 to 11 carbon atoms; and

c) mixtures of a) and b).

2. A bleach activator according to Claim 1 wherein R^1 is selected from the group consisting of heptyl, octyl, nonyl, decyl, undecyl, decenyl, 2,4,4-trimethylpentyl, 1-ethylpentyl, and mixtures thereof.
3. A bleach activator according to Claim 2 selected from the group consisting of nonanoyl valerolactam, isononanoyl valerolactam and 2-ethylhexanoyl valerolactam.
4. A bleach activator according to Claim 1 wherein R^2 , R^3 , R^4 , and R^5 are H and R^6 is selected from the group consisting of H, methyl, methoxy, ethyl,

ethoxy, propyl, propoxy, isopropyl, isopropoxy, butyl, tert-butyl, butoxy, tert-butoxy, pentyl, pentoxy, hexyl, hexoxy, Cl, and NO₂.

5. A bleach activator according to Claim 4 selected from the group consisting of benzoyl valerolactam, 3-chlorobenzoyl valerolactam and 4-nitrobenzoyl valerolactam.
6. A bleach activator according to Claim 1 wherein R², R³, R⁴ are H, and R⁵ and R⁶ are members selected from the group consisting of methyl, methoxy, and Cl.
7. A laundry bleaching system comprising:
 - i) at least about 0.1% by weight of a peroxygen bleaching compound; and
 - ii) at least about 0.1% by weight of an acyl valerolactam bleach activator according to Claim 1.
8. A bleaching system according to Claim 7 wherein said peroxygen bleaching compound is percarbonate and said valerolactam activator is selected from the group consisting of benzoyl valerolactam, nonanoyl valerolactam, isononanoyl valerolactam, 2-ethylhexanoyl valerolactam, 3-chlorobenzoyl valerolactam, and 4-nitrobenzoyl valerolactam.
9. A laundry detergent composition comprising at least about 0.1% of the bleaching system according to Claim 7 and at least about 1% by weight of conventional detergent ingredients.
10. A laundry detergent composition according to Claim 9 wherein said conventional detergent ingredients comprise a protease enzyme.
11. A composition according to Claim 9 wherein said conventional detergent ingredients comprise from about 5% to about 80% of a deterative surfactant; said peroxygen bleaching compound is percarbonate or perborate.
12. A composition according to Claim 11 further comprising a second bleach activator selected from the group consisting of tetraacetyl ethylene diamine, alkanoyloxybenzenesulfonate, benzoxazin-type activators, peroxy-

- 5 acid activators having amide moieties, N-acyl caprolactams and mixtures thereof.
13. A composition according to Claim 12 wherein R¹ is selected from the group consisting of heptyl, octyl, nonyl, decyl, undecyl, decenyl, 2,4,4-trimethylpentyl, 1-ethylpentyl, and mixtures thereof; and the second bleach
5 activator is tetraacetyl ethylene diamine.
14. A composition according to Claim 12 wherein R², R³, R⁴, and R⁵ are H and R⁶ is selected from the group consisting of H, methyl, methoxy, ethyl, ethoxy, propyl, propoxy, isopropyl, isopropoxy, butyl, tert-butyl, butoxy,
5 tert- butoxy, pentyl, pentoxy, hexyl, hexoxy, Cl, and NO₂; and the second bleach activator is tetraacetyl ethylene diamine.
15. A method for cleaning fabrics, said method comprising contacting said fabrics in an aqueous liquor comprising a bleaching system which comprises:
5 a) at least about 0.1% by weight of a peroxygen bleaching compound capable of yielding hydrogen peroxide in an aqueous solution; and
b) at least about 0.1% by weight of one or more acyl valerolactam bleach activators according to Claim 1.
16. A method according to Claim 15 wherein the peroxygen bleaching compound is selected from the group consisting of sodium perborate monohydrate, sodium perborate tetrahydrate, sodium pyrophosphate peroxyhydrate, urea peroxyhydrate, sodium percarbonate, sodium peroxide
5 and mixtures thereof.
17. A method according to Claim 16 wherein said bleaching system further comprises a second bleach activator.
18. A method according to Claim 17 wherein said second bleach activator is selected from the group consisting of alkanoyloxybenzenesulfonate, tetraacetyl ethylene diamine, benzoxazin-type activators, peroxyacid
5 activators having an amide moiety, N-acyl caprolactams and mixtures thereof.

- 5 19. A method for cleaning fabrics in automatic washing machines having parts made of natural rubber which is susceptible to oxidative degradation, said method comprising agitating said fabrics in said machine in an aqueous liquor comprising a bleaching system according to Claim 7 such that said natural rubber parts of said machine are substantially undamaged by the bleaching system.
20. A liquid bleach composition comprising:
- 5 a) from about 1% to about 25% by weight of a bleach activator according to Claim 1; and
- b) from about 0.1% to about 10% by weight of a peroxygen bleaching compound comprising hydrogen peroxide or which is capable of yielding hydrogen peroxide in an aqueous solution.
21. A liquid bleach composition according to Claim 20 further comprising from about 1% to about 20% of a phase stabilizer.

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 94/06798

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 5 C11D3/39 C07D211/76

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,3 206 439 (F.E. DETORO ET AL.) 14 September 1965 see example 7 ---	1,5
X	JOURNAL OF ORGANIC CHEMISTRY, vol.52, no.8, 1987 pages 1599 - 1601 R. MOORE ET AL 'New 2-Pyridone Derivates via 1,3-Dipolar Cycloadditions of novel Mesoionic Compounds' ---	1
X	CHEM. PHARM. BULL., vol.34, no.3, 1986 pages 975 - 979 M. MASUI ET AL. 'Anodic Oxidation of Amides and Lactams using N-Hydroxyphthalimide as a Mediator' --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- 'a' document member of the same patent family

Date of the actual completion of the international search

2 November 1994

Date of mailing of the international search report

11.11.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Grittern, A

INTERNATIONAL SEARCH REPORT

Inter- national Application No

PCT/US 94/06798

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF ORGANIC CHEMISTRY, vol.42, no.10, 1977 pages 18081808 - 1809 'N- vs. C-Acylation of Metalated O-Methylactims' ---	1
X	CHEMICAL ABSTRACTS, vol. 94, no. 5, 2 February 1981, Columbus, Ohio, US; abstract no. 29760, page 487 ; see abstract & COLLECT. CZECH. CHEM. COMMUN., vol.45, no.9, 1980 pages 2524 - 2531 J. STEHLICEK ET AL. 'Aminolysis of N-acyl lactams' ---	1
X	CHEMICAL ABSTRACTS, vol. 108, no. 22, 30 May 1988, Columbus, Ohio, US; abstract no. 187402, page 8 ; see abstract & CS,A,239 425 (J. STEHLICEK ET AL.) ---	1
A	DATABASE WPI Derwent Publications Ltd., London, GB; AN 89-211471 & JP,A,1 149 726 (TAKI CHEMICAL KK) 12 June 1989 see abstract ---	1
A	US,A,4 248 928 (G.L. SPADINI ET AL.) 3 February 1981 see column 6, line 3 - line 27 -----	1,7

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 94/06798

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-3206439		NONE	
CS-A-239425		NONE	
US-A-4248928	03-02-81	GB-A- 1586769	25-03-81
		AT-B- 374822	12-06-84
		CA-A- 1118964	02-03-82
		CH-A- 633041	15-11-82
		DE-A,C 2744642	13-04-78
		FR-A,B 2367114	05-05-78
		JP-C- 1475292	18-01-89
		JP-A- 53091917	12-08-78
		JP-B- 63011396	14-03-88
		NL-A- 7710964	10-04-78
		SE-B- 432264	26-03-84
		SE-A- 7711151	07-04-78
		US-A- 4179390	18-12-79
		BE-A- 859461	06-04-78